Tumor Board Tuesday – Dr. Hossein Borghaei, 08/30/2022: Stage III Unresectable NSCLC

Posttest Rationale

- 1. What adjuvant treatment would you select for a patient with unresectable stage III NSCLC initially treated with definitive chemoradiation?
 - a. Chemotherapy
 - b. Durvalumab
 - c. Observation
 - d. Osimertinib

Rationale: Durvalumab is FDA indicated for the treatment of adult patients with unresectable, stage III NSCLC whose disease has not progressed following concurrent platinum-based chemotherapy and radiation therapy. Neither chemotherapy, observation, nor osimertinib are appropriate adjuvant treatment options in this scenario.

References: IMFINZI® (durvalumab) [prescribing information]. AstraZeneca. Approved 2017. Revised July 2021. https://www.accessdata.fda.gov/drugsatfda docs/label/2021/761069s028lbl.pdf

National Comprehensive Cancer Network (NCCN). NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines®): Non-Small Cell Lung Cancer (v3.2022). Updated March 16, 2022. Accessed April 13, 2022. https://www.nccn.org/professionals/physician_gls/pdf/nscl.pdf

- 2. What treatment regimen would you select for a patient with unresectable stage 3 NSCLC and 36% PD-L1 expression who received adjuvant durvalumab treatment?
 - e. Atezolizumab + chemo
 - f. Nivolumab + ipilimumab
 - g. Nivolumab + ipilimumab + chemo
 - h. Pembrolizumab + chemo

Rationale: Pembrolizumab + pemetrexed + (carboplatin or cisplatin) is the NCCN guideline-preferred treatment regimen for this patient; the other options are listed as "other recommended" or "useful in certain circumstances".

Reference: National Comprehensive Cancer Network (NCCN). NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines®): Non-Small Cell Lung Cancer (v3.2022). Updated March 16, 2022. Accessed April 13, 2022. https://www.nccn.org/professionals/physician_gls/pdf/nscl.pdf